

Secondary metabolites from *Scabiosa atropurpurea* and their antioxidant and xanthine oxidase inhibitory activities

Ceren ÖZTÜRK^{1*} , Nisa Beril ŞEN¹ , Etil GÜZELMERİÇ¹ , Hasan KIRMIZİBEKMEZ¹ 

¹ Department of Pharmacognosy, Faculty of Pharmacy, Yeditepe University, Kayışdağı, İstanbul, Türkiye

* Corresponding Author. E-mail:ceren.ozturk@yeditepe.edu.tr(C.Ö.); Tel. +90-539-624 98 88.

Received: 31 March 2023/ Revised: 15 June 2023 / Accepted: 19 June 2023

ABSTRACT: Two secoiridoid glycosides, Ionicajaposide I (1) and secologanin dimethyl acetal (2), six flavonoids, isoorientin (3) hesperidin (4), apigenin 7-*O*-neohesperidoside (5), luteolin 7-*O*-β-D-glucopyranoside (6), kaempferol 3-*O*-(3'',6''-di-(*E*)-*p*-coumaroyl)-β-D-glucopyranoside (7), and kaempferol 3-*O*-(3''-*O*-acetyl, 6''-*O*-(*E*)-*p*-coumaroyl)-β-D-glucopyranoside (8), two caffeoylquinic acid derivatives, chlorogenic acid (9) and 3,5-dicaffeoylquinic acid (10) were isolated from the EtOAc and H₂O subextracts of the crude MeOH extract prepared from the aerial parts of *Scabiosa atropurpurea*. Their structures were identified by extensive 1D and 2D NMR experiments as well as ESI-MS analysis. Xanthine oxidase (XO) inhibitory and antioxidant activities of the isolated compounds were evaluated by *in vitro* tests. Compounds, 3, 5, 6, 9 and 10 showed mild-to-moderate inhibitory effects on XO enzyme. The highest antioxidant activity was found for compound 10 according to results of DPPH, FRAP and CUPRAC assays. This is the first study on the XO inhibitory activities for compounds 1, 2, 5, 7 and 8.

KEYWORDS: *Scabiosa atropurpurea*; secoiridoids; flavonoids; caffeoylquinic acid derivatives; xanthine oxidase inhibitory activity; antioxidant activity

1. INTRODUCTION

The genus *Scabiosa* belongs to Caprifoliaceae family (formerly Dipsacaceae) and comprises approximately 100 species worldwide, mainly distributed in the Mediterranean region [1,2]. Various species belonging to this genus have been utilized in different folk medicines for their potential benefits on human health particularly against pulmonary and cutaneous diseases [3]. For instance, *Scabiosa succisa* is recommended for the treatment of diphtheria and respiratory infections including bronchitis, influenza, bronchial pneumonia as well as asthma. It is also employed as a remedy for skin disorders such as ulcer and certain types of dermatoses like scabies or herpes ringworm [4-6]. The genus *Scabiosa* is represented by 30 species in the flora of Türkiye [7]. *S. atropurpurea* L., a biennial or perennial plant growing wild in Türkiye, is known as "Moruyuzotu" [8]. This species is orally consumed to regulate menstrual cycle in Northern Peru while it is used to treat measles and furuncles in Catalonia. A herbal tea obtained from the aerial parts of *S. atropurpurea* is indicated as a veterinary diuretic in Iberian Peninsula, whereas an infusion of its flowers is applied externally for acne [1,8-10]. Besides, it is preferred as an ornamental plant due to its bluish-lilac flowers [11].

Biological studies on *S. atropurpurea* extracts showed different pharmacological effects including antimicrobial, antioxidant, antihyperglycemic and hepatoprotective [1,10-12]. Previous phytochemical investigations on this species revealed the presence of iridoids and phenolic compounds [11-13]. However, there is no detailed study on the isolation of its secondary metabolites and their *in vitro* xanthine oxidase (XO) inhibitory and antioxidant activities.

The excess production of uric acid in the serum leads to hyperuricemia which further increases the risk of gout and related metabolic disorders [14,15]. Xanthine oxidase is a critical enzyme in uric acid synthesis that catalyses the oxidation of hypoxanthine to xanthine and further, xanthine to uric acid with reactive oxygen species (ROS) production [16]. Therefore, the inhibition of XO activity will help reduce the development of hyperuricemia and gout by interrupting the transformation of xanthine to uric acid and thereby reducing uric acid levels in the bloodstream as well as contributing to antioxidant activity by decreasing ROS generation [17,18]. Although allopurinol, a well-known synthetic XO inhibitor, is clinically used for the management of

How to cite this article: Öztürk C, Şen NB, Güzelmeriç E, Kirmızıbekmez H. Secondary metabolites from *S. atropurpurea* and their antioxidant and xanthine oxidase inhibitory activities. J Res Pharm. 2024; 28(1): 79-88.

hyperuricemia and gout, it may cause some side effects like gastrointestinal distress, renal toxicity or allergic reactions. Hence, it is important to develop a more safer XO inhibitor for the treatment of gout and hyperuricemia [19,20].

In present study, we aimed to isolate the secondary metabolites from the aerial parts of *S. atropurpurea* L. by using chromatographic techniques. Further, antioxidant and xanthine oxidase inhibitory activities of the isolates were evaluated by *in vitro* methods as some flavonoids were previously reported to inhibit XO enzyme significantly [21,22].




Figure 1. Chemical structures of compounds 1-10 isolated from *S. atropurpurea*.

2. RESULTS

2.1. Structure elucidation of the isolates

The air-dried and powdered aerial parts of *S. atropurpurea* were extracted with MeOH. The crude MeOH extract was suspended in H₂O and submitted to liquid-liquid extraction with CHCl₃ and EtOAc, respectively. Then organic solvents were evaporated to dryness to yield CHCl₃, EtOAc and remaining H₂O subextracts. EtOAc and H₂O subextracts were subjected to various chromatographic separations to afford ten secondary metabolites including two secoiridoid glycosides (1 and 2), six flavonoids (3-8) and two caffeoylquinic acid derivatives (9 and 10) (Figure 1). The isolates were identified as lonicejaposide I (1) [23], secologanin dimethyl acetal (2) [24], isoorientin (3) [25], hesperidin (4) [26], apigenin 7-O-neohesperidoside (5) [27], luteolin 7-O- β -D-glucopyranoside (6) [28], kaempferol 3-O-(3'',6''-di-(E)-p-coumaroyl)- β -D-glucopyranoside (7) [29], kaempferol 3-O-(3''-O-acetyl, 6''-O-(E)-p-coumaroyl)- β -D-glucopyranoside (8) [30], chlorogenic acid (9) and 3,5-dicaffeoylquinic acid (10) [31] by comparing their spectroscopic data with literature values. In previous studies, compounds 6 and 10 were characterized by LC-MS analyses from the extracts of *S. atropurpurea* while compounds 5 and 9 were recently detected in this species [10,11,32]. Previous studies on different *Scabiosa* species revealed the presence of apigenin, luteolin and kaempferol derivatives as the main flavonoids which

is similar with our results [1]. Among the flavonoid glycosides herein reported, compounds **3**, **4**, **7** and **8** are reported from *S. atropurpurea* for the first time. Iridoids and secoiridoids are widely distributed in Caprifoliaceae family, especially in the genus *Scabiosa* [33]. The occurrence of secoiridoids were previously reported in *S. atropurpurea*, but compounds **1** and **2** are being reported for the first time from this species [13]. Moreover, compounds **1**, **2**, and **4** are new for the genus *Scabiosa*. Kaempferol 3-O-(3''-O-acetyl, 6''-O-(*E*)-*p*-coumaroyl)- β -D-glucopyranoside (**8**) is an acetylated kaempferol derivative first isolated from *Scabiosa hymettia*. Structurally, it is similar to *trans*-tiliroside, another kaempferol derivative with a 6-O-(*p*-coumaroyl)- β -D-glucopyranosyl moiety [30]. Its chemical structure was elucidated on the basis of detailed 1D and 2D NMR techniques (Figure 2-5) as well as MS analysis. To our knowledge, it is being reported for the third time from the genus *Scabiosa* after *S. hymettia* and *S. stellata*. Moreover, it was only shown to possess *in vitro* antimicrobial activity [30,34].




Figure 2. ^1H NMR Spectrum (400 MHz, CD_3OD) of **8**.




Figure 3. ^{13}C NMR Spectrum (100 MHz, CD_3OD) of **8**.




Figure 4. HSQC Spectrum of 8.




Figure 5. HMBC Spectrum of 8.

2.2. Antioxidant activity

Antioxidant effects of compounds (**1-10**) were assessed *in vitro* by DPPH, FRAP and CUPRAC methods (Table 1). The results of antioxidant tests were expressed as mg of Trolox equivalents (TE) per gram of isolated compounds (mg TE/g isolated compound). Amongst the tested isolates, 3,5-dicaffeoylquinic acid (**10**) showed the highest antioxidant activity in DPPH, FRAP and CUPRAC assays. Lonicejasposide I (**1**), a secoiridoid derivative, displayed the lowest antioxidant activity in FRAP and CUPRAC methods while possessed no activity in DPPH test.

Generally, the number of free phenolic hydroxyl groups on the chemical structures of natural compounds are directly proportional with their antioxidant activities [35]. A number of researchers have reported that caffeoylquinic acid derivatives possess a high antioxidant effects. Indeed, the antioxidant capacity of caffeoylquinic acid derivatives depend on the presence of caffeoyl units in the molecule [36,37]. Thus, compound **10** (dicaffeoylquinic acid derivative) exhibited higher antioxidant activity than compound **9** (monocaffeoylquinic acid derivative). Regarding flavonoids, the glycosylation on their phenolic hydroxyl groups diminishes their antioxidant activity. In our study, isoorientin (**3**) displayed better antioxidant activity than hesperidin (**4**), apigenin 7-*O*-neohesperidoside (**5**) and luteolin 7-*O*- β -D-glucopyranoside (**7**) whose C-7 (OH) groups were substituted with sugar units. In contrast to flavonoids, caffeoylquinic acids are not glycosylated at their phenolic hydroxyl groups [38]. According to our results, flavonoid glycosides (**3-8**) showed lower antioxidant effect than caffeoylquinic acid derivatives (**9** and **10**) possibly due to their glycosylation at C-3 or C-7 positions. Previous studies demonstrated that iridoids and secoiridoids did not exhibit a strong antioxidant activity as similar with our findings. This is probably owing to their non-phenolic structures [39,40].

Table 1. Antioxidant activity results of compounds **1-10** in DPPH, FRAP, CUPRAC assays.

Compound	DPPH	FRAP mg TE/g	CUPRAC
1	N.A.	24.62 $\pm$ 0.44 ^g	34.16 $\pm$ 0.81 ^h
2	N.A.	116.93 $\pm$ 2.69 ^d	25.31 $\pm$ 0.95 ^h
3	718.94 $\pm$ 1.20 ^b	896.09 $\pm$ 17.34 ^b	330.28 $\pm$ 19.52 ^c
4	12.84 $\pm$ 1.53 ^c	402.51 $\pm$ 6.36 ^c	201.83 $\pm$ 0.58 ^d
5	N.A.	46.05 $\pm$ 1.06 ^{fg}	78.84 $\pm$ 2.28 ^g
6	10.34 $\pm$ 2.87 ^c	66.99 $\pm$ 0.59 ^{ef}	140.69 $\pm$ 2.00 ^f
7	N.A.	123.40 $\pm$ 14.99 ^d	185.13 $\pm$ 0.85 ^{de}
8	N.A.	93.04 $\pm$ 7.98 ^{de}	165.17 $\pm$ 1.03 ^e
9	720.39 $\pm$ 3.02 ^b	879.84 $\pm$ 3.54 ^b	362.39 $\pm$ 12.14 ^b
10	748.20 $\pm$ 7.24 ^a	1038.49 $\pm$ 27.14 ^a	640.59 $\pm$ 7.75 ^a

DPPH: 2,2-diphenyl-1-picrylhydrazyl; FRAP: Ferric-reducing antioxidant power; CUPRAC: Cupric-reducing antioxidant capacity; N.A.: Not active. All experiments were done in thrice and results were expressed as means $\pm$ standard deviation (SD). The means with different superscript letters in each column (^{a-h}) demonstrate no significant differences ($p < 0.05$).

2.3. Xanthine oxidase inhibitory activity

Xanthine oxidase plays a key role in the formation of uric acid, the end product of purine metabolism. This enzyme catalyses the oxidative hydroxylation of hypoxanthine and xanthine to produce uric acid. The high concentration of uric acid in serum results in hyperuricemia which is a risk factor for gout [41,42]. Therefore, inhibitors of XO decrease serum uric acid levels and are potentially useful for the treatment of gout associated with hyperuricemia [43]. In our study, compounds (**1-10**) isolated from the *S. atropurpurea* were evaluated for their XO inhibitory activities (Table 2).

Among the tested compounds, isoorientin (**3**), apigenin 7-*O*-neohesperidoside (**5**), luteolin 7-*O*- β -D-glucopyranoside (**6**), chlorogenic acid (**9**) and 3,5-dicaffeoylquinic acid (**10**) displayed mild-to-moderate XO inhibitory activities, with IC₅₀ values ranging from 9.58 to 48.67 μ g/mL. In particular, luteolin 7-*O*- β -D-glucopyranoside (**6**) showed the highest XO inhibitory activity (IC₅₀: 9.58 $\pm$ 1.98 μ g/mL) which is comparable to positive control, allopurinol (IC₅₀: 2.13 $\pm$ 0.10 μ g/mL).

Previously, luteolin 7-*O*- β -D-glucopyranoside (**6**) was reported to have mild-to-moderate XO inhibitory effect with an IC₅₀ value of 26.55 μ M, while allopurinol showed activity with IC₅₀: 0.92 μ M [44]. On the other hand, Sarawek et al. found out that **6** possessed moderate XO inhibitory activity (IC₅₀: 19.90 μ M) which was five times less potent than reference drug, allopurinol (IC₅₀: 3.65 μ M) [45]. Similar results for luteolin 7-*O*- β -D-glucopyranoside (IC₅₀: 9.58 $\pm$ 1.98 μ g/mL) were observed in our study (allopurinol, IC₅₀: 2.13 $\pm$ 0.10 μ g/mL).

Isoorientin displayed weak XO inhibitory effect (IC_{50} : $117.2 \pm 13.5 \mu M$) compared to positive control, quercetin (IC_{50} : $7.7 \pm 0.3 \mu M$) [46]. An et al. also reported that isoorientin exerted 96% inhibition of XO enzyme at concentration of $667 \mu mol/L$ which is similar to allopurinol at $80 \mu mol/L$ concentration with 97% inhibitory ratio [47]. This study also proved that isoorientin showed weak inhibitory activity against XO which seems to be compatible with our results. Chlorogenic acid (**9**) and 3,5-dicaffeoylquinic acid (**10**) were found to inhibit XO enzyme with an IC_{50} value of $28.29 \mu M$ and $43.86 \mu M$, respectively (allopurinol, IC_{50} : $2.49 \mu M$) [48]. However, in our study, **10** showed higher XO inhibitory activity than **9**. Previous *in vitro* experiment demonstrated that hesperidin (**4**) exhibited relatively very weak inhibitory effect (IC_{50} : $>2000 mg/L$) against XO which is line with our finding (did not show activity in the tested concentration ranges, $50-400 \mu g/mL$) [49]. The lack of the activity of the tested kaempferol derivatives (**7** and **8**) may be related to their glycosidic structures, since their aglycone, kaempferol, was found to significantly inhibit the XO enzyme [50]. Yuan et al. also reported that the glycosylation of some flavonoids, such as kaempferol, might cause to decrease in their inhibition toward XO enzyme [51]. This is the first study to evaluate XO inhibitory activities of compounds **1**, **2**, **5**, **7** and **8**.

Table 2. *In vitro* xanthine oxidase (XO) inhibitory activities of compounds (**1-10**).

Compound	XO Inhibitory Activity (IC_{50} , $\mu g/mL \pm SD$)
1	N.A.
2	N.A.
3	48.67 ± 6.46^c
4	N.A.
5	42.58 ± 3.29^c
6	9.58 ± 1.98^a
7	N.A.
8	N.A.
9	27.32 ± 2.40^b
10	22.38 ± 2.42^b
Allopurinol	2.13 ± 0.10

N.A.: Not active. The IC_{50} values of each sample were measured in triplicate. Results were stated as means $\pm$ SD. The means with lowercase letters (^{a-c}) indicate no significant differences ($p < 0.05$).

3. CONCLUSION

Phytochemical studies on the EtOAc and H_2O subextracts of *S. atropurpurea* yielded 10 secondary metabolites belonging to secoiridoid glycosides (**1** and **2**), flavonoids (**3-8**) and caffeoylquinic acid derivatives (**9** and **10**). Their structures were determined based on the spectroscopic methods. To our best knowledge, compounds **1**, **2**, and **4** are being purified and identified for the first time from the genus *Scabiosa*. Compounds **3**, **5**, **6**, **9** and **10** showed mild-to-moderate inhibitory activity on the XO enzyme, compared to the positive control. In particular, luteolin 7- O - β -D-glucopyranoside (**6**) displayed the highest inhibition against XO enzyme and 3,5-dicaffeoylquinic acid possessed the highest antioxidant activity among all tested compounds.

4. MATERIALS AND METHODS

4.1. General experimental procedures

Column chromatography (CC) was performed on silica gel 60 (Merck, Darmstadt, Germany), Sephadex LH-20 (Sigma-Aldrich, St. Louis, MO, USA) and Polyamide (Fluka Analytical, Sigma-Aldrich, USA). Medium-Pressure Liquid Chromatography (MPLC) was performed with Sepacore® Flash Systems X10/X50 (BuchiLabortechnik AG, Flawil, Switzerland) on RediSep columns (LiChroprep C₁₈: 150 and 100 g; SiO₂: 12 g, Teledyne Isco, Lincoln, Nebraska, USA). Thin Layer Chromatography (TLC) analyses were carried out on silica gel 60 F₂₅₄ plates (Merck, Darmstadt, Germany), visualization was accomplished by spraying with 1% vanillin/ H_2SO_4 followed by heating at $105^\circ C$ for 2-3 min, and detected with UV lights (254 and 365 nm). Analytical grade solvents (CH_2Cl_2 and MeOH) were used for chromatographic separations. 1D (1H : 400 MHz, ^{13}C : 100 MHz) and 2D NMR (COSY, HSQC, HMBC and ROESY) spectra were recorded by using a Varian Mercury FT spectrometer (Palo Alto, USA) in CD_3OD or $DMSO-d_6$. TMS was used as an internal standard. The chemical shift values (δ) were expressed in ppm while coupling constants (J) were in Hz. ESI-MS data was recorded on Agilent G6530B TOF/Q-YTOF mass spectrometer (Agilent Technologies, USA) in positive ion mode.

4.2. Plant material

The aerial parts of *Scabiosa atropurpurea* L. were collected from Yeditepe University, Kayışdağı campus, İstanbul province of Türkiye in July 2021. The plant material was identified by Prof. Dr. Hasan Kırmızıbekmez. A voucher specimen (YEF 21005) has been stored in the Herbarium of the Department of Pharmacognosy, Faculty of Pharmacy, Yeditepe University, İstanbul, Türkiye.

4.3. Extraction and isolation

The air-dried and powdered aerial parts of *S. atropurpurea* (250 g) were extracted two times with MeOH (2.5 L) at 45 °C for 2 h to obtain crude MeOH extract (40 g, yield: 16%) after removal of solvent under vacuum. An aliquot of MeOH extract (39.5 g) was dispersed in H₂O (80 mL) and then partitioned with CHCl₃ (3 x 80 mL) and EtOAc (3 x 80 mL), respectively to yield CHCl₃ (9.69 g), EtOAc (1.57 g), and rH₂O (25.42 g) subextracts. EtOAc subextract (1.57 g) was applied to C₁₈-MPLC (150 g) with a gradient of H₂O-MeOH (0-100 % MeOH) to give 16 fractions (Frs. A-P). Fr. N (42 mg) was further separated by silica gel (SiO₂) (8 g) CC using CH₂Cl₂-MeOH as a solvent system (98:2 → 90:10) to obtain **2** (12 mg). Purification of **7** (10 mg) was accomplished by separation of fr. P (37 mg) on SiO₂ (6 g) column with CH₂Cl₂-MeOH mixture (95:5 → 90:10). Fr. O (21 mg) was subjected to SiO₂ (5 g) CC with CH₂Cl₂-MeOH as the eluent (96:4 → 90:10) to yield **8** (3 mg). Fr. J (84 mg) was chromatographed over SiO₂ (15 g) CC by elution of CH₂Cl₂-MeOH-H₂O mixture (95:5:0 → 80:20:2) to give six subfractions, frs. J₁-J₆. Fr. J₅ (20 mg) was further purified by Sephadex LH-20 (22 g) CC using MeOH to afford **6** (6 mg). Fr. Compound **10** (9 mg) was isolated from fr. G (105 mg) by SiO₂ (15 g) CC with CH₂Cl₂-MeOH-H₂O as the eluting solvent (97:3:0 → 70:30:3). H₂O subextract (25.42 g) was fractionated over polyamide column (100 g) eluting with stepwise gradient of MeOH in H₂O (0-100%, in steps of 20%, each 200 mL) to obtain eight fractions (Frs. 1-8). Fr. 2 (5.841 g) was further fractionated on SiO₂ (120 g) column using CH₂Cl₂-MeOH-H₂O (90:10:1 → 61:32:7) as mobile phase to give eight subfractions, frs. 2a-h. Fr. 2d (565 mg) was implemented to Sephadex LH-20 (95 g) CC eluted with MeOH to give four fractions, frs. 2d₁₋₄. Purification of **4** (3 mg) was carried out from fr. 2d₄ (27 mg) via SiO₂-MPLC (12 g) by gradient elution of MeOH in CH₂Cl₂ (95:5 to 80:20). Fr. 2d₃ (444 mg) was applied to C₁₈-MPLC (100 g) with a gradient of H₂O-MeOH mixture (20 to 60% MeOH) and obtained **1** (16 mg). Fr. 6 (556 mg) was subjected to C₁₈-MPLC (150 g) using H₂O-MeOH (linear gradient 10-50%) as the eluting solvent to give **9** (49 mg) and together with seven fractions, frs. 6a-g. Among these fractions, fr. 6c (12 mg) was separated over Sephadex LH-20 (6 g) CC eluted with MeOH to yield **3** (6 mg). Purification of fr. 6g (39 mg) over SiO₂ (8 g) CC using CH₂Cl₂-MeOH-H₂O (95:5:0 → 85:15:1) mixtures afforded **5** (3 mg). Compound **3** (15 mg) was purified from fr. 6d (103 mg) by SiO₂ (15 g) CC eluting with CH₂Cl₂-MeOH-H₂O (95:5:0 → 80:20:2). Fr. 8 (50 mg) was chromatographed on Sephadex LH-20 (22 g) using MeOH as the mobile phase and obtained **9** (14 mg) together with 1 fraction, fr. 8a. Fr. 8a (22 mg) was further loaded in column that contains 4 g silica gel eluted with CH₂Cl₂-MeOH (90:10 → 85:15) to give **6** (12 mg).

4.4. Antioxidant activity

4.4.1. Preparation of standard solution for antioxidant assays

The stock standard solution of Trolox (Sigma-Aldrich, Steinheim, Germany) was prepared in MeOH and further diluted to prepare working standard solutions in the range of from 200 µg/mL to 3.125 µg/mL before DPPH, FRAP and CUPRAC analyses.

4.4.2. DPPH method

DPPH assay was performed according to the method of Degirmencioglu et al. [52] with some modifications. 20 µL of working concentrations of Trolox standard solutions, sample test solution or or blank (MeOH) were added into a 96-well plate, separately. Thereafter, 280 µL of 0.1 mM methanolic DPPH solution (abs. ~0.7, Sigma-Aldrich, Steinheim, Germany) was inserted into each well resulting the final volume of 300 µL. The plate was incubated at room temperature for 30 min in the dark conditions. Lastly, the absorbance was measured at 520 nm using the plate reader Varioskan™ LUX (Thermo Fisher Scientific, USA). Results were expressed as mg of Trolox equivalent (TE)/g.

4.4.3. FRAP method

Firstly, FRAP solution was freshly prepared by mixing 1 volume of iron (III) chloride solution (2x10⁻² M, Riedel-de Haen, Germany), 1 volume of TPTZ solution (1x10⁻² M), and 10 volumes of sodium acetate buffer (pH 3.6) solution. Then, 280 µL of FRAP solution and 20 µL of sample test solution or Trolox standard solution

or blank (water) were added into a 96-well microplate. After 6 minutes, the absorbance was read spectrophotometrically at 593 nm [53]. Outcomes of the analysis were stated as mg TE/g.

4.4.4. CUPRAC method

Evaluation of the antioxidant activities of the compounds by the CUPRAC method were carried out according to study of Apak et al. [54]. Initially, 85 µL of 10 mM Copper (II) sulfate pentahydrate (Sigma-Aldrich, Steinheim, Germany) was mixed with 85 µL of 7.5 mM neocuproine, 85 µL ammonium acetate buffer solution (pH 7) and 51 µL of distilled water in wells of a 96-well microplate. Then, either 43 µL of sample test solution or Trolox standard solution or blank (water) were inserted in each well. After 30 min of kept of the plate at room temperature, the absorbance was detected at 450 nm. Findings were expressed as mg TE/g.

4.5. Xanthine oxidase inhibitory activity

The inhibitory effect on xanthine oxidase was conducted according to the procedure of Azmi et al. [16] with minor changes. 75 µL of 50 mM sodium phosphate buffer (pH 7.5), 25 µL of sample solution dissolved in water or DMSO, 25 µL of freshly prepared enzyme solution (0.2 U/mL of XO in buffer solution) and 25 µL of distilled water were added into a 96-well microplate, respectively. The reaction mixture was pre-incubated at 37°C for 15 min and followed by addition of 50 µL substrate solution (0.15 mM xanthine, Alfa Aesar, Haverhill, Massachusetts, USA) into the mixture. Then, the mixture was incubated at 37°C for 30 min, and the reaction was terminated by adding 50 µL of 0.5 M HCl. Finally, the absorbance was read at 290 nm against a blank, containing all reagents except enzyme solution. Allopurinol (Acros Organics, Geel, Belgium) used as a positive control which was prepared at different concentrations in range of from 15 to 200 µg/mL. The xanthine oxidase inhibition (%) was calculated by using the following formula:

$$\text{XO inhibition (\%)} = [1 - (\text{Sample}_{\text{abs}} - \text{Sample blank}_{\text{abs}}) / (\text{Control}_{\text{abs}} - \text{Control blank}_{\text{abs}})] \times 100$$

4.6. Statistical analysis

All experiments were performed in triplicate. Experimental results and statistical significance of the results were determined by using Minitab 17 software with one-way analysis of variance (ANOVA). Comparison of significance was ascertained by Tukey post hoc-test with 95% confidence. Results were expressed as means $\pm$ SD and significance levels were defined as $p < 0.05$.

Author contributions: Concept – C.Ö., H.K.; Design – C.Ö., H.K.; Supervision – H.K.; Resources – C.Ö., H.K.; Materials – C.Ö., H.K.; Data Collection and/or Processing – C.Ö., H.K., N.B.Ş., E.G.; Analysis and/or Interpretation – C.Ö., H.K., N.B.Ş., E.G.; Literature Search – C.Ö., H.K., N.B.Ş., E.G.; Writing – C.Ö., H.K., N.B.Ş., E.G.; Critical Reviews – C.Ö., H.K., N.B.Ş., E.G.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] Pinto DCGA, Rahmouni N, Beghidja N, Silva AMS. *Scabiosa* Genus: A rich source of bioactive metabolites. *Medicines*. 2018; 5(4): 110. <https://doi.org/10.3390/medicines5040110>
- [2] Lehbili M, AlabdulMagid A, Hubert J, Kabouche A, Voutquenne-Nazabadioko L, Renault JH, Nuzillard JM, Morjani H, Abedini A, Gangloff SC, Kabouche Z. Two new bis-iridoids isolated from *Scabiosa stellata* and their antibacterial, antioxidant, anti-tyrosinase and cytotoxic activities. *Fitoterapia*. 2018; 125: 41-48. <https://doi.org/10.1016/j.fitote.2017.12.018>
- [3] Bendamene S, Boutaghane N, Bellik Y, Sayagh C, AlabdukMagid A, Harakat D, Kabouche Z, Voutquenne-Nazabadioko L. Semipapposides A-M, triterpenoid bidesmosides saponins from the roots of *Scabiosa semipapposa*. *Phytochemistry*. 2020; 180: 112526. <https://doi.org/10.1016/j.phytochem.2020.112526>
- [4] BesbesHilila M, Mosbah H, Majouli K, Ben Nejma A, Ben Jannet H, Mastouri M, Aouni M, Selmi B. Antimicrobial activity of *Scabiosa arenaria*Forssk. extracts and pure compounds using bioguided fractionation. *Chem Biodivers*. 2016; 13(10): 1262-1272. <https://doi.org/10.1002/cbdv.201600028>
- [5] Al-Qudah MA, Ootom NK, Al-Jaber HI, Saleh AM, Abu Zarga MH, Afifi FU, Abu Orabi ST. New flavonol glycoside from *Scabiosa prolifera* L. aerial parts with in vitro antioxidant and cytotoxic activities. *Nat Prod Res*. 2017; 31(24): 2865-2874. <http://doi.org/10.1080/14786419.2017.1305377>
- [6] Kılınç H, Masullo M, D'Urso G, Karayildirim T, Alankus O, Piacente S. Phytochemical investigation of *Scabiosa sicula* guided by a preliminary HPLC-ESIMSⁿ profiling. *Phytochemistry*. 2020; 174: 112350. <https://doi.org/10.1016/j.phytochem.2020.112350>

- [7] Matthews VA. *Scabiosa* L. In: Davis PH. (Ed). Flora of Turkey and East Aegean Islands. Edinburgh University Press, Edinburgh, 1972; pp.602-621.
- [8] Erarslan ZB, Yeşil Y. The anatomical properties of *Scabiosa atropurpurea* L. (Caprifoliaceae). Istanbul J Pharm. 2018; 48(1): 1-5. <https://doi.org/10.5152/IstanbulJPharm.2018.376278>
- [9] Jalloul AB, Garzoli S, Chaar H, elJribi C, Abderrabba M. Seasonal effect on bioactive compounds recovery using aqueous extraction, antioxidant activities, and volatile profiles of different parts of *S. maritima* (*Scabiosa atropurpurea* sub. *maritima* L.). S Afr J Bot. 2023; 152: 63-79. <https://doi.org/10.1016/j.sajb.2022.11.031>
- [10] Hrichi S, Chaabane-Banaoues R, Bayar S, Flamini G, Oulad El Majdoub Y, Mangraviti D, Mondello L, El Mzoughi R, Babba H, Mighri Z, Cacciola F. Botanical and genetic identification followed by investigation of chemical composition and biological activities on the *Scabiosa atropurpurea* L. stem from Tunisian flora. Molecules. 2020; 25(21): 5032. <http://doi.org/10.3390/molecules25215032>
- [11] Kilinc H, Masullo M, Lauro G, D'Urso G, Alankus O, Bifulco G, Piacente S. *Scabiosa atropurpurea*: A rich source of iridoids with α -glucosidase inhibitory activity evaluated by in vitro and in silico studies. Phytochemistry. 2023; 205: 113471. <https://doi.org/10.1016/j.phytochem.2022.113471>
- [12] Elhawary SS, Eltantawy ME, Sleem AA, Abdallah HM, Mohamed NM. Investigation of phenolic content and biological activities of *Scabiosa atropurpurea* L. World ApplSci J. 2011; 15(3): 311-317.
- [13] Polat E, Alankus-Caliskan Ö, Karayildirim T, Bedir E. Iridoids from *Scabiosa atropurpurea* L. subsp. *maritima* Arc. (L.). Biochem Syst Ecol 2010; 38(2): 253-255. <https://doi.org/10.1016/j.bse.2010.01.004>
- [14] Umamaheswari M, AsokKumar K, Somasundaram A, Sivashanmugam T, Subhadradevi V, Ravi TK. Xanthine oxidase inhibitory activity of some Indian medical plants. J Ethnopharmacol. 2007; 109(3): 547-551. <https://doi.org/10.1016/j.jep.2006.08.020>
- [15] Chen J, Li Q, Ye Y, Ran M, Ruan Z, Jin N. Inhibition of xanthine oxidase by theaflavin: Possible mechanism for anti-hyperuricaemia effect in mice. Process Biochem. 2020; 97: 11-18. <https://doi.org/10.1016/j.procbio.2020.06.024>
- [16] Azmi SM, Jamal P, Amid A. Xanthine oxidase inhibitory activity from potential Malaysian medicinal plant as remedies for gout. Int Food Res J. 2012; 19(1): 159-165.
- [17] Song JU, Jang JW, Kim TH, Park H, Park WS, Jung SH, Kim GT. Structure-based design and biological evaluation of novel 2-(indol-2-yl) thiazole derivatives as xanthine oxidase inhibitors. Bioorg Med Chem Lett. 2016; 26(3): 950-954. <https://doi.org/10.1016/j.bmcl.2015.12.055>
- [18] Mohamed Isa SSP, Ablat A, Mohamad J. The Antioxidant and xanthine oxidase inhibitory activity of *Plumeria rubra* flowers. Molecules. 2018; 23(2): 400. <https://doi.org/10.3390/molecules23020400>
- [19] Bardin T. Current management of gout in patients unresponsive or allergic to allopurinol. Jt Bone Spine. 2004; 71(6): 481-485. <https://doi.org/10.1016/j.jbspin.2004.07.006>
- [20] Pacher P, Nivorozhkin A, Szabó C. Therapeutic effects of xanthine oxidase inhibitors: renaissance half a century after the discovery of allopurinol. Pharmacol Rev. 2006; 58(1): 87-114. <https://doi.org/10.1124/pr.58.1.6>
- [21] Lin S, Zhang G, Liao Y, Pan J, Gong D. Dietary flavonoids as xanthine oxidase inhibitors: Structure–affinity and structure–activity relationships. J Agric Food Chem. 2015; 63(35): 7784-7794. <https://doi.org/10.1021/acs.jafc.5b03386>
- [22] Cos P, Ying L, Calomme M, Hu JP, Cimanga K, Van Poel B, Pieters L, Vlietinck AJ, Berghe DV. Structure–activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. J Nat Prod. 1998; 61(1): 71-76. <https://doi.org/10.1021/np970237h>
- [23] Wan J, Jiang CX, Tang Y, Ma GL, Tong YP, Jin ZX, Zang Y, Osman EE, Li J, Xiong J, Hu JF. Structurally diverse glycosides of secoiridoid, bisiridoid, and triterpene-bisiridoid conjugates from the flower buds of two Caprifoliaceae plants and their ATP-citrate lyase inhibitory activities. Bioorg Chem. 2022; 120: 105630. <https://doi.org/10.1016/j.bioorg.2022.105630>
- [24] Kawai H, Kuroyanagi M, Ueno A. Iridoid glucosides from *Lonicera japonica* THUNB. Chemical Pharm Bull. 1988; 36(9): 3664-3666. <https://doi.org/10.1248/cpb.36.3664>
- [25] Çaliş I, Birincioğlu SS, Kırmızıbekmez H, Pfeiffer B, Heilmann J. Secondary metabolites from *Asphodelus aestivus*. ZNaturforsch B. 2006; 61(10): 1304-1310. <https://doi.org/10.1515/znb-2006-1019>
- [26] KadioğluYaman B, Şen O, Salman A, Sipahi H, Kúsz N, Hohmann J, Kırmizimekbez H. Anti-inflammatory effects of *Lyciumbarbarum* leaf extracts in lipopolysaccharide-induced RAW 264.7 macrophage cells and isolation of secondary metabolites. J Res Pharm. 2019; 23(4): 740-748. <https://doi.org/10.12991/jrp.2019.183>
- [27] Zhang H, Li X, Wu K, Wang M, Liu P, Wang X, Deng R. Antioxidant activities and chemical constituents of flavonoids from the flower of *Paeonia ostii*. Molecules. 2016; 22(1): 5. <https://doi.org/10.3390/molecules22010005>
- [28] Akdemir ZS, Tatlı II, Bedir E, Khan, IA Antioxidant flavonoids from *Verbascum salviifolium*Boiss. Fabad J Pharm Sci. 2003; 28: 71-75.
- [29] Liu H, Orjala J, Sticher O, Rali T. Acylated flavonol glycosides from leaves of *Stenochlaena palustris*. J Nat Prod. 1999; 62(1): 70-75. <https://doi.org/10.1021/np980179f>
- [30] Christopoulou C, Graikou K, Chinou I. Chemosystematic value of chemical constituents from *Scabiosa hymettia* (Dipsacaceae). ChemBiodivers. 2008; 5(2): 318-323. <https://doi.org/10.1002/cbdv.200890029>
- [31] Indy Tamayose C, Dos Santos EA, Roque N, Costa-Lotufo LV, Pena Ferreira MJ. Caffeoylquinic acids: Separation method, antiradical properties and cytotoxicity. ChemBiodivers. 2019; 16(7): e1900093. <https://doi.org/10.1002/cbdv.201900093>

- [32] Ben Toumia I, Sobeh M, Ponassi M, Banelli B, Dameriha A, Wink M, ChekirGhedira L, Rosano C. A methanol extract of *Scabiosa atropurpurea* enhances doxorubicin cytotoxicity against resistant colorectal cancer cells in vitro. *Molecules*. 2020; 25(22): 5265. <https://doi.org/10.3390/molecules25225265>
- [33] Papalexandrou A, Magiatis P, Perdetzoglou D, Skaltsounis AL, Chinou IB, Harvala C. Iridoids from *Scabiosa variifolia* (Dipsacaceae) growing in Greece. *Biochem Syst Ecol*. 2003; 1(31): 91-93.
- [34] Lehbili M, AlabdulMagid A, Kabouche A, Voutquenne-Nazabadioko L, Morjani H, Harakat D, Kabouche Z. Triterpenoid saponins from *Scabiosa stellata* collected in North-eastern Algeria. *Phytochemistry*. 2018; 150: 40-49. <https://doi.org/10.1016/j.phytochem.2018.03.005>
- [35] Olszowy M. What is responsible for antioxidant properties of polyphenolic compounds from plants?. *Plant Physiol Biochem*. 2019; 144: 135-143. <https://doi.org/10.1016/j.plaphy.2019.09.039>
- [36] Jiang XW, Bai JP, Zhang Q, Hu XL, Tian X, Zhu J, Liu J, Meng WH, Zhao QC. Caffeoylquinic acid derivatives from the roots of *Arctium lappa* L.(burdock) and their structure-activity relationships (SARs) of free radical scavenging activities. *Phytochem Lett*. 2016; 15: 159-163. <http://doi.org/10.1016/j.phytol.2015.12.008>
- [37] Zhao JG, Yan QQ, Xue RY, Zhang J, Zhang YQ. Isolation and identification of colourless caffeoyl compounds in purple sweet potato by HPLC-DAD-ESI/MS and their antioxidant activities. *Food Chem*. 2014; 161: 22-26. <https://doi.org/10.1016/j.foodchem.2014.03.079>
- [38] Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med*. 1996; 20(7): 933-956. [https://doi.org/10.1016/0891-5849\(95\)02227-9](https://doi.org/10.1016/0891-5849(95)02227-9)
- [39] Pacifico S, D'Abrosca B, Pascarella MT, Letizia M, Uzzo P, Piscopo V, Fiorentino A. Antioxidant efficacy of iridoid and phenylethanoid glycosides from the medicinal plant *Teucrium chamaedris* in cell-free systems. *Bioorg Med Chem*. 2009; 17(17): 6173-6179. <https://doi.org/10.1016/j.bmc.2009.07.065>
- [40] Huang SW, Qiao JW, Sun X, Gao PY, Li LZ, Liu QB, Sun B, Wu DL, Song SJ. Secoiridoids and lignans from the leaves of *Diospyros kaki* Thunb. with antioxidant and neuroprotective activities. *J Funct Food*. 2016; 24:183-195. <https://doi.org/10.1016/j.jff.2016.03.025>
- [41] Wang Y, Zhang G, Pan J, Gong D. Novel insights into the inhibitory mechanism of kaempferol on xanthine oxidase. *J Agric Food*. 2015; 63(2): 526-534. <https://doi.org/10.1021/jf505584m>
- [42] Mehmood A, Ishaq M, Zhao L, Safdar B, Rehman AU, Munir M, Raza A, Nadeem M, Iqbal W, Wang C. Natural compounds with xanthine oxidase inhibitory activity: A review. *Chem Biol Drug Des*. 2019; 93(4): 387-418. <https://doi.org/10.1111/cbdd.13437>
- [43] Li Q, Shi C, Wang M, Zhou M, Liang M, Zhang T, Yuan E, Wang Z, Yao M, Ren J. Tryptophan residue enhances in vitro walnut protein-derived peptides exerting xanthine oxidase inhibition and antioxidant activities. *J Funct Foods*. 2019; 53: 276-285. <https://doi.org/10.1016/j.jff.2018.11.024>
- [44] Qu L, Ruan JY, Jin LJ, Shi WZ, Li XX, Han LF, Zhang Y, Wang T. Xanthine oxidase inhibitory effects of the constituents of *Chrysanthemum morifolium* stems. *Phytochem Lett*. 2017; 19: 39-45. <http://doi.org/10.1016/j.phytol.2016.11.007>
- [45] Sarawek S, Feistel B, Pischel I, Butterweck V. Flavonoids of *Cynara scolymus* possess potent xanthinoxidase inhibitory activity in vitro but are devoid of hypouricemic effects in rats after oral application. *Planta Med*. 2008; 74(03): 221-227. <https://doi.org/10.1055/s-2008-1034316>
- [46] Pham AT, Nguyen C, Malterud KE, Diallo D, Wangenstein H. Bioactive flavone-C-glycosides of the African medicinal plant *Biophytum umbellatum*. *Molecules*. 2013; 18(9): 10312-10319. <https://doi.org/10.3390/molecules180910312>
- [47] An MF, Wang MY, Shen C, Sun ZR, Zhao YL, Wang XJ, Sheng J. Isoorientin exerts a urate-lowering effect through inhibition of xanthine oxidase and regulation of the TLR4-NLRP3 inflammasome signaling pathway. *J Nat Med*. 2021; 75(1): 129-141. <https://doi.org/10.1007/s11418-020-01464-z>
- [48] Li Y, Wan Y, Li R, Xu L, Xie M, Fu G. Solvent extraction of caffeoylquinic acids from *Artemisia selengensis* Turcz leaves and their in vitro inhibitory activities on xanthine oxidase. *Ind Crops Prod*. 2018; 118: 296-301. <https://doi.org/10.1016/j.indcrop.2018.03.055>
- [49] Kim DS, Lim SB. Semi-continuous subcritical water extraction of flavonoids from *Citrus unshiu* peel: Their antioxidant and enzyme inhibitory activities. *Antioxidants*. 2020; 9(5): 360. <https://doi.org/10.3390/antiox9050360>
- [50] Kostić DA, Dimitrijević DS, Stojanović GS, Palić IR, Đorđević AS, Ickovski JD. Xanthine oxidase: Isolation, assays of activity, and inhibition. *J Chem*. 2015; 2015(2): 1-8. <https://doi.org/10.1155/2015/294858>
- [51] Yuan M, Liu Y, Xiao A, Leng J, Liao L, Ma L, Liu L. The interaction of dietary flavonoids with xanthine oxidase in vitro: Molecular property-binding affinity relationship aspects. *RSC Adv*. 2019; 9(19): 10781-10788. <https://doi.org/10.1039/c8ra09926j>
- [52] Degirmencioglu HT, Guzelmeric E, Yuksel PI, Kirmizibekmez H, Deniz I, Yesilada E. A new type of Anatolian propolis: Evaluation of its chemical composition, activity profile and botanical origin. *Chem Biodivers*. 2019; 16(12): e1900492. <https://doi.org/10.1002/cbdv.201900492>
- [53] Berker KI, Güçlü K, Demirata B, Apak R. A novel antioxidant assay of ferric reducing capacity measurement using ferrozine as the colour forming complexation reagent. *Anal Methods*. 2010; 2(11): 1770-1778. <https://doi.org/10.1039/C0AY00245C>
- [54] Apak R, Güçlü K, Özyürek M, Karademir SE. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J Agric Food Chem*. 2004; 52(26): 7970-7081. <https://doi.org/10.1021/jf048741x>